

Published in final edited form as:

*J Cardiovasc Transl Res.* 2014 March ; 7(2): 203–211. doi:10.1007/s12265-013-9515-7.

## Coagulation, Protease Activated Receptors and Viral Myocarditis

**Silvio Antoniak and Nigel Mackman**

Department of Medicine, Division of Hematology and Oncology, UNC McAllister Heart Institute, University of North Carolina at Chapel Hill, 98 Manning Drive Campus Box 7035, Chapel Hill, North Carolina, USA

### Abstract

The coagulation protease cascade plays an essential role in hemostasis. In addition, a clot contributes to host defense by limiting the spread of pathogens. Coagulation proteases induce intracellular signaling by cleavage of cell surface receptors called protease-activated receptors (PARs). These receptors allow cells to sense changes in the extracellular environment, such as infection. Viruses activate the coagulation cascade by inducing tissue factor expression and by disrupting the endothelium. Virus infection of the heart can cause myocarditis, cardiac remodeling and heart failure. Recent studies using a mouse model have shown that tissue factor, thrombin and PAR-1 signaling all positively regulate the innate immune during viral myocarditis. In contrast, PAR-2 signaling was found to inhibit interferon- $\beta$  expression and the innate immune response. These observations suggest that anticoagulants may impair the innate immune response to viral infection and that inhibition of PAR-2 may be a new target to reduce viral myocarditis..

### Keywords

tissue factor; protease activated receptor; thrombin; myocarditis; coxsackievirus B3; virus infection; innate immune response

---

During infection the immune system is activated to eliminate the pathogen. The coagulation system functions as part of the immune response by forming a clot that limits dissemination of pathogens. Importantly, there is crosstalk between coagulation and inflammation. In this review, we discuss recent findings on the contribution of activation of coagulation and the clotting cascade and protease activated receptors to early responses to viral infections particularly viral myocarditis.

### Myocarditis

Myocarditis is defined as inflammation of the heart muscle which can be caused by different insults, such as infections, autoimmune reactions, toxins and adverse drug reactions. Infections with enteroviruses, adenoviruses and other viruses can cause viral myocarditis. One of the first viruses identified to cause viral myocarditis was coxsackievirus B3 (CVB3) [1,2]. Viral myocarditis can be divided in three phases [3,4]. The early phase (day 0–4 post infection) is defined as early virus replication in various organs, such as pancreas, liver and heart. During this phase cardiac injury is caused by the virus itself. In the acute phase of myocarditis (days 7 to 10 post infection) there is infiltration of immune cells into the heart and increased cytokine/chemokine expression that controls viral replication. At this stage

---

Address for Correspondence: Silvio Antoniak, PhD, University of North Carolina at Chapel Hill, 98 Manning Drive Campus Box 7035, Chapel Hill, North Carolina, 27599, Telephone: (919) 843 3961, Fax : (919) 966 7639, antoniak@email.unc.edu.

**COMPETING INTERESTS STATEMENT:** None

cardiac injury is mediated by immune cells, such as auto-reactive T cells. In the last phase (after day 15 post infection) the virus is eliminated and inflammation subsides. However, viral genomes can persist in the heart. Complications of viral myocarditis can occur in early and late stages of the disease and include pathologic remodeling that can lead to impaired heart function, dilated cardiomyopathy (DCM) and heart failure [5,6].

## Innate immune response in myocarditis

The innate immune response is a first line defense system to infection by pathogens. Toll-like receptors (TLRs) are a family of innate immune receptors that play a central role in host defense by recognizing pathogen-associated molecular patterns (PAMPs) [7]. Viral infections are detected by different receptors, including the endosomal receptor TLR3. TLR3 is activated by dsRNA from dsRNA viruses and replication of ssRNA viruses. The importance of the different innate and adaptive immune responses in viral myocarditis has been reviewed by others [8–10]. Studies with mice found that increased virus-specific IgGs are detectable from day 7 onwards post CVB3 infection [11]. This indicates that innate immunity has a major role in antiviral defense [10].

Experimental myocarditis studies in mice infected with either CVB3 or CVB4 have shown a role of TLR3 and its adapter Toll/IL-1 receptor/resistance (TIR)-domain-containing adapter-inducing interferon- $\beta$  (TRIF) [12–16]. Activation of TLR3 leads to the expression of interferons (IFNs), which play a crucial role in the early innate immune response to viruses [17]. IFNs are divided into type I (IFN- $\alpha$  and IFN- $\beta$ ), type II (IFN- $\gamma$ ) and type III (IFN- $\lambda$ ). Type I, but not type II, IFN signaling has been shown to be required for an innate immune response to CVB3 infection [18,19]. Importantly, administration of IFN- $\beta$  reduces CVB3-induced myocarditis in mice and humans [16,20–22]. IFN- $\alpha$  and - $\beta$  are expressed at low levels by numerous cell types but expression is increased in virally infected cells. IFNs stimulate the expression of IFN-stimulated genes (ISGs) that help eliminate the virus [23]. The ISG 2'-5' oligoadenylate synthetase (OAS)/RNase L inhibits CVB3 replication by cleavage of viral ssRNA [18]. Other ISGs are chemokines, such as CXCL10, and these recruit natural killer (NK) cells to the site of infection [24]. Notably, CXCL10 deficiency in mice is associated with increased cardiac injury after infection [12,24]. In viral myocarditis, NK cells are one of the first cell types infiltrating the heart to limit viral replication and spread [25].

## Blood Coagulation

Tissue factor (TF) is the primary initiator of coagulation protease cascade [26]. It is the cellular receptor and cofactor for factor VII/VIIa (FVII/VIIa). TF is expressed by cells within the vessel wall and is essential for hemostasis [27,28]. It is also expressed by activated monocytes and this response is thought to be part of the innate immune response to limit dissemination of pathogens [29,30]. In addition, the clotting system activates participates in host defense. For instance, fibrin(ogen) engagement of the integrin Mac1 on macrophages is required for efficient clearance of *Staphylococcus aureus* in mice [31]. Furthermore, fibrin(ogen) modulates natural killer cell activity [32].

## Activation of coagulation in viral infections

Viral infections lead to the induction of TF expression in various cell types that are exposed to blood and this leads to activation of the coagulation cascade [4,33–37]. Interestingly, the dsRNA analog, polyinosinic-polycytidylic acid (poly IC), is a synthetic ligand for TLR3 and when administered to mice induces TF expression in endothelial cells and a hypercoagulable state [38]. Zare et al. observed that TF expression in murine splenocytes was increased by ssRNA (poly I) but not dsRNA (poly IC) [39]. This induction was

MyD88-dependent and IFN-independent [39]. Furthermore, type I and III IFN expression was associated with activation of coagulation [40]. However, IFN- $\alpha$  administration reduced platelet reactivity and ventricular thrombus formation during CVB3 myocarditis [21,41]. Furthermore, proinflammatory cytokines produced during viral infections may increase TF expression in endothelial cells and monocytes [4,35,42].

## Activation of coagulation in myocarditis

Heart failure and DCM induced by viral myocarditis is associated with increased risk of thrombosis [43]. Clinical case reports have described an increased incidence of ventricular fibrin deposition and thrombi during acute myocarditis [44–49]. The presence of ventricular thrombi during myocardial inflammation may be due to TF expression by the endocardium [46,50]. We and others have observed thrombosis in the heart in mouse models of myocarditis [4,21,37,43,51,52]. One study reported increased TF expression in the endocardium of patients with myocardial inflammation that was associated with immune cells, most likely CD45<sup>+</sup> leukocytes (activated T cells) [53]. Consistent with this, we observed an association between TF expression and markers of activated endothelial cells in heart biopsies of patients with suspected myocarditis [4]. Furthermore, we showed that CVB3 infection induces myocardial TF expression [4,37]. Our data suggest an acute but no longer lasting inflammation of the heart leads to increased TF expression that triggers thrombosis [4,42,43,54]. In the chronic phase of heart failure, other parameters, such as platelet reactivity and blood stasis, may contribute to the enhanced risk for thrombotic events, as seen in patients with chronic DCM [4,44]. It has been suggested that anti-thrombotics should be used in patients with active cardiac inflammation caused by myocarditis [42,46,55–58].

## Protease-activated receptors

There is cross-talk between the coagulation cascade and inflammatory response that is mediated, in part, by protease activated receptors (PARs) [59]. PARs belong to the family of G protein-coupled receptors [59]. There are four PARs (PAR-1 to -4). Cleavage of the N-terminal sequence of the receptor reveals a new N-terminal sequence that acts as tethered ligand by binding to the receptor and inducing its activation. A wide range of serine proteases, including immune cell-derived proteases and coagulation proteases can either activate or inactivate PARs which is dependent on cell type and post-translational modification of the receptor (Table 1). Thrombin is the primary activator of PAR-1, 3 and 4 whereas tryptase and trypsin are the primary activators of PAR-2 [60,61]. PARs are expressed by many cell types in the body and their expression patterns are comparable in different species. However, one important species specific difference in PAR expression exists; human platelets express PAR1 and PAR4 whereas mouse platelet express PAR3 and PAR4 [62]. The role of PARs in different diseases and immune responses has been reviewed by others [63,64]. PAR activation can be either pro-inflammatory or anti-inflammatory responses depending on the activator and cell type [63,64]. Moretti et al. proposed a dual-sensor system whereby an infection is detected by TLRs recognizing PAMPs and PARs being activated by extracellular proteases [65]. TLR-dependent signaling was affected by PAR expression/activation and *vice versa* [65].

## Serine proteases in viral myocarditis

Besides coagulation proteases, other serine protease are expressed and released by stromal and circulating immune cells during infection [64]. Inhibitors of serine proteinases are routinely used to reduce activation of coagulation and inflammation in the clinic, such as during cardiac surgery. One of these inhibitors is the naturally occurring proteinase inhibitor

aprotinin. Administration of aprotinin reduced thrombin generation and PAR-1 dependent signaling [66]. In addition, trypsin is ectopically expressed by cardiac cells after ssRNA virus infection [67,68]. Aprotinin reduced trypsin-mediated signaling and viral replication in a model of influenza A myocarditis [67,68]. Trypsin stimulation of cells inhibited IFN responses after influenza A infection [69], which may be via PAR-2 activation. In addition, trypsin may increase the infectivity of influenza A via the cleavage of the hemagglutinin protein [70]. Aprotinin also reduced inflammation by preventing trypsin activation of PAR-4 and recruitment of neutrophils [71]. The known inhibitory action of aprotinin against trypsin as well as thrombin suggests that several pathways may be affected by aprotinin during virus infection.

Neutrophils secrete several proteases, such as cathepsin G, which activate PAR-1 and -4 [72]. Neutrophils are involved in clearing CVB3 from the heart [73]. Further, neutrophils contribute to T cell mediated pathologic responses in the heart during autoimmune myocarditis [74]. Cathepsin G and neutrophil elastase inactivate PAR-2 on human lung epithelial cells [75]. In addition, elastase can activate PAR-1 on epithelial cells [76]. The neutrophil elastase inhibitor ZD0892 was shown to reduce viral myocarditis by inhibiting immune cell infiltration without effecting the viral load [77]. Other proteases, such as matrix metalloproteinases (MMP), have been implicated in viral myocarditis progression [78]. MMP1 and MMP13 are able to activate PAR-1 [37,79,80]. Cardiac cells express MMP13 [79]. We found that a pan-MMP inhibitor or an MMP13 inhibitor reduced TLR3 dependent expression of CXCL10 by cardiac fibroblasts *in vitro* (Figure) [37]. In addition, line with these findings, MMP13 inhibition increased the virus levels and cardiac injury in CVB3 infected mice [37].

Mast cells provide another rich source of serine proteases, such as tryptase and chymase, which are activators of PAR-2 [81,82]. Mast cells degranulate within six hours of CVB3 infection [83]. Furthermore, mast cell deficient mice develop significantly less myocarditis than control littermates when infected with encephalomyocarditis virus [84]. However, the effect of inhibitors of mast cell tryptase, such as by protamine [85], and other commercially available tryptase inhibitors, such as APC366 [86], has not been studied.

Interestingly, the virus genome itself can encode proteases [87]. Enterovirus protease 3C was shown to exhibit papain/chymase-like activity and inhibition of enterovirus protease 3C reduced virus levels [88]. Papain can stimulate PAR-2 dependent signaling in T-cells [89]. Furthermore, papain-like viral proteases blocked dsRNA-induced IFN- $\beta$  expression by interfering with IRF3 activation [90]. In addition, chymase pretreatment of fibroblasts rapidly inhibited the ability of these cells to respond to thrombin [91]. The data suggest that virally encoded papain/chymase-like proteases may activate PAR-2 and inactivate PAR-1 and thus reduce anti-viral innate immune responses.

## PARs and viral myocarditis

PAR-1 is expressed by cardiomyocytes, cardiac fibroblasts, smooth muscle cells, endothelial cells and leukocytes [60]. During viral infection, thrombin mediated PAR-1 activation was linked to increased susceptibility to herpes viruses of endothelial cells *in vitro* [92]. We recently investigated the role of PAR-1 in a mouse CVB3 myocarditis model. Early after infection, we observed reduced expression of IFN- $\beta$  and the IFN response gene CXCL10 as well as NK cell infiltration into the heart in PAR-1 deficient mice compared to wild-type mice [37]. These changes were associated with a significant increase in viral load in the heart and increased cardiac injury 8 days post infection [37]. Using bone-marrow transplantation, we found that PAR-1 on non-hematopoietic cells played a major role in the immune response [37]. *In vitro* studies revealed that PAR-1 stimulation on cardiac

fibroblasts enhanced TLR3-dependent p38 activation and expression of IFN- $\beta$  and CXCL10 (Figure) [37]. Further, PAR-1 activation was associated with increased NK cell activity of murine splenocytes [37]. This is in agreement with an earlier study showing NK cell activity can be increased by thrombin [93].

PAR-2 is expressed by cardiomyocytes, cardiac fibroblasts, smooth muscle cells, endothelial cells and immune cells [60,64]. TF-dependent PAR-2 activation increased herpes simplex virus infection of endothelial cells *in vitro* [94]. In collaboration with Rauch's group we found that PAR-2 deficient mice were protected from CVB3 myocarditis compared to infected wild type mice [95]. An absence of PAR-2 was associated with reduced viral load in the heart and myocarditis. Interestingly, hearts and isolated cardiac fibroblasts of PAR-2 deficient mice expressed higher levels of IFN- $\beta$  [95]. *In vitro* experiments revealed that PAR-2 activation reduced TLR3-dependent expression of IFN- $\beta$  and IFN-responses genes in cardiac fibroblasts as well as epithelial cells [95,96]. This observation may be due to a direct interaction of PAR-2 with TLR3 which subsequently leads to inhibition of TLR3 signaling [95]. Others showed that PAR-2 contains a TIR domain that may interact with adaptor proteins, such as MyD88 and TRIF [97]. In contrast to the results with TLR3, PAR-2 enhances TLR4 dependent signaling [97,98]. With regard to myocarditis, TLR4 was shown to contribute to disease progression [99–102]. PAR-2 expression was positively correlated with myocardial inflammation and negatively with IFN- $\beta$  expression and heart function in hearts of patients with non-ischemic cardiomyopathy [95]. Furthermore, Pan et al. showed that PAR-2 inhibition reduced influenza replication in cardiomyocytes *in vitro* [68].

PAR-4 is known as the main signaling receptor for thrombin on mouse platelets [62]. Platelets play a central role in hemostasis and thrombosis [103]. However, they also contribute to other processes, such as inflammation and immunity [104,105]. PAR-4 is also expressed by cardiomyocytes and smooth muscle cells in the heart as well as circulating immune cells [60,64,106]. It was proposed that PAR-4 mediates a pro-survival signal within the heart and that a lack of PAR-4 increases apoptosis and loss of cardiomyocytes [60]. The role of PAR-4 in infections has not been investigated in detail. However, we observed increased plasma cardiac troponin I levels in PAR-4 deficient mice compared to wild-type mice 8 days after CVB3 infection (Antoniak and Mackman, unpublished data). It is not clear whether these findings were due to activation of PAR-4 on platelets or other cells. Interestingly, non-steroid anti-inflammatory drugs (NSAIDs) including aspirin, decrease platelet reactivity. Mice studies and case reports suggested that the use of NSAIDs is associated with increased mortality in virus infections including myocarditis [107–110].

## Anticoagulation

There are few studies reporting the effect of anticoagulation on viral myocarditis. Treatment with low molecular weight heparin begun either before or after virus inoculation reduced mortality after CVB3 infection of mice [57]. Heparin reduced collagen deposition and fibrosis in later stages of myocarditis but did not affect viral replication. However, heparin increased cardiac inflammation in the acute phase of infection [57]. The mechanism of this effect is not certain but may be due to reduced fibrin deposition and fibrosis [56]. Importantly, heparin also has anti-adhesive activity that may explain some of the effects. We observed increased virus levels and cardiac injury 8 days after CVB3 infection of mice treated with either a TF inhibitor or thrombin inhibitor compared to control treated mice [37]. These results suggested that TF mediated thrombin generation enhances antiviral responses that limit cardiac injury.

Schnitt et al. have suggested the use of a FXIII inhibitor or a fibrinolytic in viral myocarditis to reduce deposition of cross-linked fibrin and fibrin-mediated fibrosis [56]. However, our



results in a mouse model suggested that upstream coagulation proteases, such as thrombin, play a protective role in viral myocarditis [37]. One important question is does the use of anticoagulants increase the susceptibility and severity of viral myocarditis? Warfarin has been the main stay for long-term anticoagulation for many years. More recently, the thrombin inhibitor dabigatran etexilate and the factor Xa inhibitor rivaroxaban have been approved for long-term anticoagulation [111]. While the risk of thrombosis is reduced in patients our results with a mouse model suggest that reduced thrombin activity may increase the susceptibility to viral infection [37]. In response to our publication [37], investigators from the RE-LY study stated that “Pradaxa® (dabigatran etexilate) did not increase the incidence of virus infections when compared to warfarin” [112]. The authors further noted that other anticoagulants, such as warfarin or FXa inhibitors, should also increase the susceptibility of virus infections [112]. Rauch’s group treated wild-type mice with the FXa inhibitor fondaparinux sodium and observed increased cardiac inflammation but reduced fibrosis and improved heart function 8 days after CVB3 infection compared to the placebo treated group [113]. At present, it is unclear if anticoagulation affects the frequency and intensity of viral infection.

## Conclusions

Recent studies indicate that coagulation proteases and PARs modulate the innate immune responses to virus infections in mouse models, including myocarditis (Table 2). Interestingly, PAR-1 enhances while PAR-2 inhibits TLR3 signaling in cardiac fibroblasts. The activators of PAR-2 in myocarditis are not yet characterized. Importantly, our findings suggest that anticoagulation may increase the risk and severity of certain virus infections. Furthermore, inhibition of PAR-2 could be a new interventional strategy to treat viral myocarditis. One must be cautious in extrapolating from mice studies to humans since human myocarditis is highly diverse [114]. With regards to the effect of anticoagulation, there are no current clinical trials to determine if anticoagulants increase the incidence and severity of virus infections. One would need to compare infections in demographic, co-morbidity matched controls with or without anticoagulant therapy. Secondly, it is possible that divergent results will be observed with different anticoagulants. Further studies are needed to investigate the role of coagulation proteases, PARs and platelets in different virus infections.

## Clinical Relevance

Basic research studies have increased our understanding of the molecular mechanism of viral myocarditis and revealed promising pathways for interventions. Our data suggest a potential new treatment option for viral myocarditis is inhibition of PAR-2 [95]. Several studies has described the development of PAR-2 inhibitors that are potential candidates for clinical trials [115,116]. However, to date the new knowledge on viral myocarditis has not resulted in any substantial improvement in the treatment of viral myocarditis [114]. This is likely due to the fact that viral myocarditis is often diagnosed too late after infection. One positive note is that Kuhl and coworkers showed that IFN- $\beta$  treatment reduced virus levels and improved heart function and survival in patients with enterovirus myocarditis [20,22].

## Acknowledgments

We thank Dr. Julie C. Williams for her careful and critical review of this manuscript. This work was supported by grants from the Myocarditis Foundation (to S. Antoniak) and the National Institutes of Health (to N. Mackman, HL119523).

## References

1. Shauer A, Gotsman I, Keren A, Zwas DR, Hellman Y, Durst R, Admon D. Acute viral myocarditis: current concepts in diagnosis and treatment. *The Israel Medical Association journal : IMAJ*. 2013; 15(3):180–185. [PubMed: 23662385]
2. Schultheiss HP, Kuhl U, Cooper LT. The management of myocarditis. *European heart journal*. 2011; 32(21):2616–2625. [PubMed: 21705357]
3. Liu PP, Mason JW. Advances in the understanding of myocarditis. *Circulation*. 2001; 104(9):1076–1082. [PubMed: 11524405]
4. Antoniak S, Boltzen U, Riad A, Kallwellis-Opara A, Rohde M, Dorner A, Tschöpe C, Noutsias M, Pauschinger M, Schultheiss HP, Rauch U. Viral myocarditis and coagulopathy: increased tissue factor expression and plasma thrombogenicity. *Journal of molecular and cellular cardiology*. 2008; 45(1):118–126. [PubMed: 18495150]
5. Imazio M, Cooper LT. Management of myopericarditis. *Expert review of cardiovascular therapy*. 2013; 11(2):193–201. [PubMed: 23405840]
6. Blauwet LA, Cooper LT. Myocarditis. *Progress in cardiovascular diseases*. 2010; 52(4):274–288. [PubMed: 20109598]
7. Thompson JM, Iwasaki A. Toll-like receptors regulation of viral infection and disease. *Advanced drug delivery reviews*. 2008; 60(7):786–794. [PubMed: 18280610]
8. Kembell CC, Alirezaei M, Whitton JL. Type B coxsackieviruses and their interactions with the innate and adaptive immune systems. *Future microbiology*. 2010; 5(9):1329–1347. [PubMed: 20860480]
9. Huber S. Host immune responses to coxsackievirus B3. *Current topics in microbiology and immunology*. 2008; 323:199–221. [PubMed: 18357771]
10. Yajima T. Viral myocarditis: potential defense mechanisms within the cardiomyocyte against virus infection. *Future microbiology*. 2011; 6(5):551–566. [PubMed: 21585262]
11. Leipner C, Grun K, Schneider I, Gluck B, Sigusch HH, Stelzner A. Coxsackievirus B3-induced myocarditis: differences in the immune response of C57BL/6 and Balb/c mice. *Medical microbiology and immunology*. 2004; 193(2–3):141–147. [PubMed: 14593475]
12. Weinzierl AO, Szalay G, Wolburg H, Sauter M, Rammensee HG, Kandolf R, Stevanovic S, Klingel K. Effective chemokine secretion by dendritic cells and expansion of cross-presenting CD4<sup>−</sup>/CD8<sup>+</sup> dendritic cells define a protective phenotype in the mouse model of coxsackievirus myocarditis. *Journal of virology*. 2008; 82(16):8149–8160. [PubMed: 18550677]
13. Negishi H, Osawa T, Ogami K, Ouyang X, Sakaguchi S, Koshiba R, Yanai H, Seko Y, Shitara H, Bishop K, Yonekawa H, Tamura T, Kaisho T, Taya C, Taniguchi T, Honda K. A critical link between Toll-like receptor 3 and type II interferon signaling pathways in antiviral innate immunity. *Proceedings of the National Academy of Sciences of the United States of America*. 2008; 105(51):20446–20451. [PubMed: 19074283]
14. Richer MJ, Lavalley DJ, Shanina I, Horwitz MS. Toll-like receptor 3 signaling on macrophages is required for survival following coxsackievirus B4 infection. *PloS one*. 2009; 4(1):e4127. [PubMed: 19122812]
15. Abston ED, Coronado MJ, Bucek A, Bedja D, Shin J, Kim JB, Kim E, Gabrielson KL, Georgakopoulos D, Mitzner W, Fairweather D. Th2 regulation of viral myocarditis in mice: different roles for TLR3 versus TRIF in progression to chronic disease. *Clinical & developmental immunology*. 2012; 2012:129486. [PubMed: 22013485]
16. Riad A, Westermann D, Zietsch C, Savvatis K, Becher PM, Bereswill S, Heimesaat MM, Lettau O, Lassner D, Dorner A, Poller W, Busch M, Felix SB, Schultheiss HP, Tschöpe C. TRIF is a critical survival factor in viral cardiomyopathy. *J Immunol*. 2011; 186(4):2561–2570. [PubMed: 21239721]
17. Newman KC, Riley EM. Whatever turns you on: accessory-cell-dependent activation of NK cells by pathogens. *Nature reviews Immunology*. 2007; 7(4):279–291.
18. Wessely R, Klingel K, Knowlton KU, Kandolf R. Cardiospecific infection with coxsackievirus B3 requires intact type I interferon signaling: implications for mortality and early viral replication. *Circulation*. 2001; 103(5):756–761. [PubMed: 11156890]

19. Deonarain R, Cerullo D, Fuse K, Liu PP, Fish EN. Protective role for interferon-beta in coxsackievirus B3 infection. *Circulation*. 2004; 110(23):3540–3543. [PubMed: 15249500]
20. Kuhl U, Pauschinger M, Schwimmbeck PL, Seeberg B, Lober C, Noutsias M, Poller W, Schultheiss HP. Interferon-beta treatment eliminates cardiotropic viruses and improves left ventricular function in patients with myocardial persistence of viral genomes and left ventricular dysfunction. *Circulation*. 2003; 107(22):2793–2798. [PubMed: 12771005]
21. Wang YX, da Cunha V, Vincelette J, White K, Velichko S, Xu Y, Gross C, Fitch RM, Halks-Miller M, Larsen BR, Yajima T, Knowlton KU, Vergona R, Sullivan ME, Croze E. Antiviral and myocyte protective effects of murine interferon-beta and -{alpha}2 in coxsackievirus B3-induced myocarditis and epicarditis in Balb/c mice. *American journal of physiology Heart and circulatory physiology*. 2007; 293(1):H69–H76. [PubMed: 17434974]
22. Kuhl U, Lassner D, von Schlippenbach J, Poller W, Schultheiss HP. Interferon-Beta improves survival in enterovirus-associated cardiomyopathy. *Journal of the American College of Cardiology*. 2012; 60(14):1295–1296. [PubMed: 23017536]
23. Fensterl V, Sen GC. Interferons and viral infections. *Biofactors*. 2009; 35(1):14–20. [PubMed: 19319841]
24. Yuan J, Liu Z, Lim T, Zhang H, He J, Walker E, Shier C, Wang Y, Su Y, Sall A, McManus B, Yang D. CXCL10 inhibits viral replication through recruitment of natural killer cells in coxsackievirus B3-induced myocarditis. *Circulation research*. 2009; 104(5):628–638. [PubMed: 19168435]
25. Godeny EK, Gauntt CJ. Murine natural killer cells limit coxsackievirus B3 replication. *J Immunol*. 1987; 139(3):913–918. [PubMed: 3036947]
26. Mackman N. Role of tissue factor in hemostasis and thrombosis. *Blood cells, molecules & diseases*. 2006; 36(2):104–107.
27. Drake TA, Morrissey JH, Edgington TS. Selective cellular expression of tissue factor in human tissues. Implications for disorders of hemostasis and thrombosis. *The American journal of pathology*. 1989; 134(5):1087–1097. [PubMed: 2719077]
28. Fleck RA, Rao LV, Rapaport SI, Varki N. Localization of human tissue factor antigen by immunostaining with monospecific, polyclonal anti-human tissue factor antibody. *Thrombosis research*. 1990; 59(2):421–437. [PubMed: 2237820]
29. Degen JL, Bugge TH, Goguen JD. Fibrin and fibrinolysis in infection and host defense. *Journal of thrombosis and haemostasis : JTH* 5 Suppl. 2007; 1:24–31.
30. Esmon CT, Xu J, Lupu F. Innate immunity and coagulation. *Journal of thrombosis and haemostasis : JTH*. 2011; 9(Suppl 1):182–188. [PubMed: 21781254]
31. Flick MJ, Du X, Witte DP, Jirouskova M, Soloviev DA, Busuttill SJ, Plow EF, Degen JL. Leukocyte engagement of fibrin(ogen) via the integrin receptor alphaMbeta2/Mac-1 is critical for host inflammatory response in vivo. *The Journal of clinical investigation*. 2004; 113(11):1596–1606. [PubMed: 15173886]
32. Palumbo JS, Talmage KE, Massari JV, La Jeunesse CM, Flick MJ, Kombrinck KW, Jirouskova M, Degen JL. Platelets and fibrin(ogen) increase metastatic potential by impeding natural killer cell-mediated elimination of tumor cells. *Blood*. 2005; 105(1):178–185. [PubMed: 15367435]
33. Key NS, Vercellotti GM, Winkelmann JC, Moldow CF, Goodman JL, Esmon NL, Esmon CT, Jacob HS. Infection of vascular endothelial cells with herpes simplex virus enhances tissue factor activity and reduces thrombomodulin expression. *Proceedings of the National Academy of Sciences of the United States of America*. 1990; 87(18):7095–7099. [PubMed: 2169619]
34. Geisbert TW, Young HA, Jahrling PB, Davis KJ, Kagan E, Hensley LE. Mechanisms underlying coagulation abnormalities in ebola hemorrhagic fever: overexpression of tissue factor in primate monocytes/macrophages is a key event. *The Journal of infectious diseases*. 2003; 188(11):1618–1629. [PubMed: 14639531]
35. Goeijenbier M, van Wissen M, van de Weg C, Jong E, Gerdes VE, Meijers JC, Brandjes DP, van Gorp EC. Review: Viral infections and mechanisms of thrombosis and bleeding. *Journal of medical virology*. 2012; 84(10):1680–1696. [PubMed: 22930518]
36. Funderburg NT, Mayne E, Sieg SF, Asaad R, Jiang W, Kalinowska M, Luciano AA, Stevens W, Rodriguez B, Brenchley JM, Douek DC, Lederman MM. Increased tissue factor expression on



- circulating monocytes in chronic HIV infection: relationship to in vivo coagulation and immune activation. *Blood*. 2010; 115(2):161–167. [PubMed: 19828697]
37. Antoniak S, Owens AP 3rd, Baunacke M, Williams JC, Lee RD, Weithauser A, Sheridan PA, Malz R, Luyendyk JP, Esserman DA, Trejo J, Kirchhofer D, Blaxall BC, Pawlinski R, Beck MA, Rauch U, Mackman N. PAR-1 contributes to the innate immune response during viral infection. *The Journal of clinical investigation*. 2013; 123(3):1310–1322. [PubMed: 23391721]
  38. Shibamiya A, Hersemeyer K, Schmidt Woll T, Sedding D, Daniel JM, Bauer S, Koyama T, Preissner KT, Kanse SM. A key role for Toll-like receptor-3 in disrupting the hemostasis balance on endothelial cells. *Blood*. 2009; 113(3):714–722. [PubMed: 18971420]
  39. Zare F, Magnusson M, Mollers LN, Jin T, Tarkowski A, Bokarewa M. Single-stranded polyinosinic acid oligonucleotides trigger leukocyte production of proteins belonging to fibrinolytic and coagulation cascades. *Journal of leukocyte biology*. 2008; 84(3):741–747. [PubMed: 18562486]
  40. Kato J, Okamoto T, Motoyama H, Uchiyama R, Kirchhofer D, Van Rooijen N, Enomoto H, Nishiguchi S, Kawada N, Fujimoto J, Tsutsui H. Interferon-gamma-mediated tissue factor expression contributes to T-cell-mediated hepatitis through induction of hypercoagulation in mice. *Hepatology*. 2013; 57(1):362–372. [PubMed: 22936459]
  41. Bhattacharyya M, Karmohapatra SK, Bhattacharya G, Bhattacharya R, Sinha AK. The role of leucocytes in the acetyl salicylic acid (aspirin) induced nitric oxide synthesis in the production of interferon-alpha, a potent inhibitor of platelet aggregation and a thrombolytic agent. *Journal of thrombosis and thrombolysis*. 2009; 28(2):173–184. [PubMed: 18846318]
  42. Antoniak S, Boltzen U, Eisenreich A, Stellbaum C, Poller W, Schultheiss HP, Rauch U. Regulation of cardiomyocyte full-length tissue factor expression and microparticle release under inflammatory conditions in vitro. *Journal of thrombosis and haemostasis : JTH*. 2009; 7(5):871–878. [PubMed: 19228282]
  43. Tomioka N, Kishimoto C, Matsumori A, Kawai C. Mural thrombi in mice with acute viral myocarditis. *Japanese circulation journal*. 1985; 49(12):1277–1279. [PubMed: 3007795]
  44. Kojima J, Miyazaki S, Fujiwara H, Kumada T, Kawai C. Recurrent left ventricular mural thrombi in a patient with acute myocarditis. *Heart and vessels*. 1988; 4(2):120–122. [PubMed: 3253272]
  45. Mazza IL, Jacobs JP, Aldousany A, Chang AC, Burke RP. Video-assisted cardioscopy for left ventricular thrombectomy in a child. *The Annals of thoracic surgery*. 1998; 66(1):248–250. [PubMed: 9692476]
  46. Weikert U, Rauch U, Kuhl U, Hohmann C, Jaster M, Pauschinger M, Schwimmbeck PL, Schultheiss HP. Increased thrombocyte activation in dilated cardiomyopathy: a risk factor for development of ventricular thrombosis despite anticoagulant therapy? *Zeitschrift fur Kardiologie*. 2002; 91(5):423–429. [PubMed: 12132290]
  47. Miyamoto K, Yasuda S, Noguchi T, Tanimoto T, Kakuchi H, Morii I, Kawamura A, Nakatani T, Nonogi H, Goto Y, Miyazaki S. Fulminant myocarditis causing severe left heart failure and massive thrombus formation following cardiac tamponade: a case report. *Journal of cardiology*. 2005; 46(1):25–31. [PubMed: 16095228]
  48. Kuh JH, Seo Y. Transatrial resection of a left ventricular thrombus after acute myocarditis. *Heart and vessels*. 2005; 20(5):230–232. [PubMed: 16160906]
  49. Lin KY, Kerur B, Witmer CM, Beslow LA, Licht DJ, Ichord RN, Kaufman BD. Thrombotic events in critically ill children with myocarditis. *Cardiology in the young*. 2013:1–8.
  50. Gottdiener JS, Gay JA, VanVoorhees L, DiBianco R, Fletcher RD. Frequency and embolic potential of left ventricular thrombus in dilated cardiomyopathy: assessment by 2-dimensional echocardiography. *The American journal of cardiology*. 1983; 52(10):1281–1285. [PubMed: 6650417]
  51. Tomioka N, Kishimoto C, Matsumori A, Kawai C. Mural thrombus in experimental viral myocarditis in mice: relation between thrombosis and congestive heart failure. *Cardiovascular research*. 1986; 20(9):665–671. [PubMed: 3791356]
  52. Kishimoto C, Ochiai H, Sasayama S. Intracardiac thrombus in murine Cocksackievirus B3 myocarditis. *Heart and vessels*. 1992; 7(2):76–81. [PubMed: 1328146]


53. Nakamura Y, Nakamura K, Fukushima-Kusano K, Ohta K, Matsubara H, Hamuro T, Yutani C, Ohe T. Tissue factor expression in atrial endothelia associated with nonvalvular atrial fibrillation: possible involvement in intracardiac thrombogenesis. *Thrombosis research*. 2003; 111(3):137–142. [PubMed: 14678810]
54. Szotowski B, Goldin-Lang P, Antoniak S, Bogdanov VY, Pathirana D, Pauschinger M, Dorner A, Kuehl U, Coupland S, Nemerson Y, Hummel M, Poller W, Hetzer R, Schultheiss HP, Rauch U. Alterations in myocardial tissue factor expression and cellular localization in dilated cardiomyopathy. *Journal of the American College of Cardiology*. 2005; 45(7):1081–1089. [PubMed: 15808768]
55. Weikert U, Kuhl U, Schultheiss HP, Rauch U. Platelet activation is increased in patients with cardiomyopathy: myocardial inflammation and platelet reactivity. *Platelets*. 2002; 13(8):487–491. [PubMed: 12487782]
56. Schnitt SJ, Stillman IE, Owings DV, Kishimoto C, Dvorak HF, Abelmann WH. Myocardial fibrin deposition in experimental viral myocarditis that progresses to dilated cardiomyopathy. *Circulation research*. 1993; 72(4):914–920. [PubMed: 7680288]
57. Frizelle S, Schwarz J, Huber SA, Leslie K. Evaluation of the effects of low molecular weight heparin on inflammation and collagen deposition in chronic coxsackievirus B3-induced myocarditis in A/J mice. *The American journal of pathology*. 1992; 141(1):203–209. [PubMed: 1321562]
58. Szotowski B, Antoniak S, Poller W, Schultheiss HP, Rauch U. Procoagulant soluble tissue factor is released from endothelial cells in response to inflammatory cytokines. *Circulation research*. 2005; 96(12):1233–1239. [PubMed: 15920023]
59. Coughlin SR. Thrombin signalling and protease-activated receptors. *Nature*. 2000; 407(6801):258–264. [PubMed: 11001069]
60. Antoniak S, Pawlinski R, Mackman N. Protease-activated receptors and myocardial infarction. *IUBMB life*. 2011; 63(6):383–389. [PubMed: 21438116]
61. Soh UJ, Dorés MR, Chen B, Trejo J. Signal transduction by protease-activated receptors. *British journal of pharmacology*. 2010; 160(2):191–203. [PubMed: 20423334]
62. Kahn ML, Zheng YW, Huang W, Bigornia V, Zeng D, Moff S, Farese RV Jr, Tam C, Coughlin SR. A dual thrombin receptor system for platelet activation. *Nature*. 1998; 394(6694):690–694. [PubMed: 9716134]
63. Steinhoff M, Buddenkotte J, Shpacovitch V, Rattenholl A, Moormann C, Vergnolle N, Luger TA, Hollenberg MD. Proteinase-activated receptors: transducers of proteinase-mediated signaling in inflammation and immune response. *Endocrine reviews*. 2005; 26(1):1–43. [PubMed: 15689571]
64. Shpacovitch V, Feld M, Bunnett NW, Steinhoff M. Protease-activated receptors: novel PARTners in innate immunity. *Trends in immunology*. 2007; 28(12):541–550. [PubMed: 17977790]
65. Moretti S, Bellocchio S, Bonifazi P, Bozza S, Zelante T, Bistoni F, Romani L. The contribution of PARs to inflammation and immunity to fungi. *Mucosal immunology*. 2008; 1(2):156–168. [PubMed: 19079173]
66. Day JR, Landis RC, Taylor KM. Aprotinin and the protease-activated receptor 1 thrombin receptor: antithrombosis, inflammation, and stroke reduction. *Seminars in cardiothoracic and vascular anesthesia*. 2006; 10(2):132–142. [PubMed: 16959740]
67. Pan HY, Yamada H, Chida J, Wang S, Yano M, Yao M, Zhu J, Kido H. Up-regulation of ectopic trypsins in the myocardium by influenza A virus infection triggers acute myocarditis. *Cardiovascular research*. 2011; 89(3):595–603. [PubMed: 21084314]
68. Pan HY, Yano M, Kido H. Effects of inhibitors of Toll-like receptors, protease-activated receptor-2 signalings and trypsin on influenza A virus replication and upregulation of cellular factors in cardiomyocytes. *The journal of medical investigation : JMI*. 2011; 58(1–2):19–28. [PubMed: 21372483]
69. Seitz C, Isken B, Heynisch B, Rettkowski M, Frensing T, Reichl U. Trypsin promotes efficient influenza vaccine production in MDCK cells by interfering with the antiviral host response. *Applied microbiology and biotechnology*. 2012; 93(2):601–611. [PubMed: 21915610]

70. Sun X, Tse LV, Ferguson AD, Whittaker GR. Modifications to the hemagglutinin cleavage site control the virulence of a neurotropic H1N1 influenza virus. *Journal of virology*. 2010; 84(17): 8683–8690. [PubMed: 20554779]
71. Gomides LF, Duarte ID, Ferreira RG, Perez AC, Francischi JN, Klein A. Proteinase-activated receptor-4 plays a major role in the recruitment of neutrophils induced by trypsin or carrageenan during pleurisy in mice. *Pharmacology*. 2012; 89(5–6):275–282. [PubMed: 22517275]
72. Sambrano GR, Huang W, Faruqi T, Mahrus S, Craik C, Coughlin SR. Cathepsin G activates protease-activated receptor-4 in human platelets. *The Journal of biological chemistry*. 2000; 275(10):6819–6823. [PubMed: 10702240]
73. Fairweather D, Frisancho-Kiss S, Yusung SA, Barrett MA, Davis SE, Steele RA, Gatewood SJ, Rose NR. IL-12 protects against coxsackievirus B3-induced myocarditis by increasing IFN-gamma and macrophage and neutrophil populations in the heart. *J Immunol*. 2005; 174(1):261–269. [PubMed: 15611248]
74. Grabie N, Hsieh DT, Buono C, Westrich JR, Allen JA, Pang H, Stavrakis G, Lichtman AH. Neutrophils sustain pathogenic CD8+ T cell responses in the heart. *The American journal of pathology*. 2003; 163(6):2413–2420. [PubMed: 14633613]
75. Dulon S, Cande C, Bunnett NW, Hollenberg MD, Chignard M, Pidard D. Proteinase-activated receptor-2 and human lung epithelial cells: disarming by neutrophil serine proteinases. *American journal of respiratory cell and molecular biology*. 2003; 28(3):339–346. [PubMed: 12594060]
76. Suzuki T, Yamashita C, Zemans RL, Briones N, Van Linden A, Downey GP. Leukocyte elastase induces lung epithelial apoptosis via a PAR-1-, NF-kappaB-, and p53-dependent pathway. *American journal of respiratory cell and molecular biology*. 2009; 41(6):742–755. [PubMed: 19307610]
77. Lee JK, Zaidi SH, Liu P, Dawood F, Cheah AY, Wen WH, Saiki Y, Rabinovitch M. A serine elastase inhibitor reduces inflammation and fibrosis and preserves cardiac function after experimentally-induced murine myocarditis. *Nature medicine*. 1998; 4(12):1383–1391.
78. Marchant D, McManus BM. Matrix metalloproteinases in the pathogenesis of viral heart disease. *Trends in cardiovascular medicine*. 2009; 19(1):21–26. [PubMed: 19467450]
79. Jaffre F, Friedman AE, Hu Z, Mackman N, Blaxall BC. beta-adrenergic receptor stimulation transactivates protease-activated receptor 1 via matrix metalloproteinase 13 in cardiac cells. *Circulation*. 2012; 125(24):2993–3003. [PubMed: 22610965]
80. Yang E, Boire A, Agarwal A, Nguyen N, O'Callaghan K, Tu P, Kuliopulos A, Covic L. Blockade of PAR1 signaling with cell-penetrating pepducins inhibits Akt survival pathways in breast cancer cells and suppresses tumor survival and metastasis. *Cancer research*. 2009; 69(15):6223–6231. [PubMed: 19622769]
81. Sharma R, Prasad V, McCarthy ET, Savin VJ, Dileepan KN, Stechschulte DJ, Lianos E, Wiegmann T, Sharma M. Chymase increases glomerular albumin permeability via protease-activated receptor-2. *Molecular and cellular biochemistry*. 2007; 297(1–2):161–169. [PubMed: 17102904]
82. Molino M, Barnathan ES, Numerof R, Clark J, Dreyer M, Cumashi A, Hoxie JA, Schechter N, Woolkalis M, Brass LF. Interactions of mast cell tryptase with thrombin receptors and PAR-2. *The Journal of biological chemistry*. 1997; 272(7):4043–4049. [PubMed: 9020112]
83. Fairweather D, Frisancho-Kiss S, Gatewood S, Njoku D, Steele R, Barrett M, Rose NR. Mast cells and innate cytokines are associated with susceptibility to autoimmune heart disease following coxsackievirus B3 infection. *Autoimmunity*. 2004; 37(2):131–145. [PubMed: 15293883]
84. Higuchi H, Hara M, Yamamoto K, Miyamoto T, Kinoshita M, Yamada T, Uchiyama K, Matsumori A. Mast cells play a critical role in the pathogenesis of viral myocarditis. *Circulation*. 2008; 118(4):363–372. [PubMed: 18606918]
85. Gan X, Liu D, Huang P, Gao W, Chen X, Hei Z. Mast-cell-releasing tryptase triggers acute lung injury induced by small intestinal ischemia-reperfusion by activating PAR-2 in rats. *Inflammation*. 2012; 35(3):1144–1153. [PubMed: 22200983]
86. He S, Aslam A, Gaca MD, He Y, Buckley MG, Hollenberg MD, Walls AF. Inhibitors of tryptase as mast cell-stabilizing agents in the human airways: effects of tryptase and other agonists of

- proteinase-activated receptor 2 on histamine release. *The Journal of pharmacology and experimental therapeutics*. 2004; 309(1):119–126. [PubMed: 14722328]
87. Bazan JF, Fletterick RJ. Viral cysteine proteases are homologous to the trypsin-like family of serine proteases: structural and functional implications. *Proceedings of the National Academy of Sciences of the United States of America*. 1988; 85(21):7872–7876. [PubMed: 3186696]
  88. Thibaut HJ, De Palma AM, Neyts J. Combating enterovirus replication: state-of-the-art on antiviral research. *Biochemical pharmacology*. 2012; 83(2):185–192. [PubMed: 21889497]
  89. Liang G, Barker T, Xie Z, Charles N, Rivera J, Druey KM. Naive T cells sense the cysteine protease allergen papain through protease-activated receptor 2 and propel TH2 immunity. *The Journal of allergy and clinical immunology*. 2012; 129(5):1377–1386 e1313. [PubMed: 22460072]
  90. Xing Y, Chen J, Tu J, Zhang B, Chen X, Shi H, Baker SC, Feng L, Chen Z. The papain-like protease of porcine epidemic diarrhea virus negatively regulates type I interferon pathway by acting as a viral deubiquitinase. *The Journal of general virology*. 2013; 94(Pt 7):1554–1567. [PubMed: 23596270]
  91. Schechter NM, Brass LF, Lavker RM, Jensen PJ. Reaction of mast cell proteases tryptase and chymase with protease activated receptors (PARs) on keratinocytes and fibroblasts. *Journal of cellular physiology*. 1998; 176(2):365–373. [PubMed: 9648924]
  92. Sutherland MR, Friedman HM, Prydzial EL. Thrombin enhances herpes simplex virus infection of cells involving protease-activated receptor 1. *Journal of thrombosis and haemostasis : JTH*. 2007; 5(5):1055–1061. [PubMed: 17461934]
  93. Naldini A, Carney DH. Thrombin modulation of natural killer activity in human peripheral lymphocytes. *Cellular immunology*. 1996; 172(1):35–42. [PubMed: 8806804]
  94. Sutherland MR, Ruf W, Prydzial EL. Tissue factor and glycoprotein C on herpes simplex virus type 1 are protease-activated receptor 2 cofactors that enhance infection. *Blood*. 2012; 119(15):3638–3645. [PubMed: 22374699]
  95. Weithaeuser A, Bobbert P, Antoniak S, Boehm A, Rauch BH, Klingel K, Savvatis K, Kroemer HK, Tschope C, Stroux A, Zeichhardt H, Poller W, Mackman N, Schultheiss HP, Rauch U. Protease-activated receptor 2 regulates the Innate Immune Response to Viral Infection in a CVB3-induced Myocarditis. *Journal of the American College of Cardiology*. 2013 in press.
  96. Nhu QM, Shirey K, Teijaro JR, Farber DL, Netzel-Arnett S, Antalis TM, Fasano A, Vogel SN. Novel signaling interactions between proteinase-activated receptor 2 and Toll-like receptors in vitro and in vivo. *Mucosal immunology*. 2010; 3(1):29–39. [PubMed: 19865078]
  97. Rallabhandi P, Nhu QM, Toshchakov VY, Piao W, Medvedev AE, Hollenberg MD, Fasano A, Vogel SN. Analysis of proteinase-activated receptor 2 and TLR4 signal transduction: a novel paradigm for receptor cooperativity. *The Journal of biological chemistry*. 2008; 283(36):24314–24325. [PubMed: 18622013]
  98. Williams JC, Lee RD, Doerschuk CM, Mackman N. Effect of PAR-2 Deficiency in Mice on KC Expression after Intratracheal LPS Administration. *Journal of signal transduction*. 2011; 2011:415195. [PubMed: 22175012]
  99. Eriksson U, Ricci R, Hunziker L, Kurrer MO, Oudit GY, Watts TH, Sonderegger I, Bachmaier K, Kopf M, Penninger JM. Dendritic cell-induced autoimmune heart failure requires cooperation between adaptive and innate immunity. *Nature medicine*. 2003; 9(12):1484–1490.
  100. Fuse K, Chan G, Liu Y, Gudgeon P, Husain M, Chen M, Yeh WC, Akira S, Liu PP. Myeloid differentiation factor-88 plays a crucial role in the pathogenesis of Coxsackievirus B3-induced myocarditis and influences type I interferon production. *Circulation*. 2005; 112(15):2276–2285. [PubMed: 16216974]
  101. Frisancho-Kiss S, Davis SE, Nyland JF, Frisancho JA, Cihakova D, Barrett MA, Rose NR, Fairweather D. Cutting edge: cross-regulation by TLR4 and T cell Ig mucin-3 determines sex differences in inflammatory heart disease. *J Immunol*. 2007; 178(11):6710–6714. [PubMed: 17513715]
  102. Roberts BJ, Dragon JA, Moussawi M, Huber SA. Sex-specific signaling through Toll-Like Receptors 2 and 4 contributes to survival outcome of Coxsackievirus B3 infection in C57Bl/6 mice. *Biology of sex differences*. 2012; 3(1):25. [PubMed: 23241283]

103. Cimmino G, Golino P. Platelet biology and receptor pathways. *Journal of cardiovascular translational research*. 2013; 6(3):299–309. [PubMed: 23307175]
104. Smyth SS, McEver RP, Weyrich AS, Morrell CN, Hoffman MR, Arepally GM, French PA, Dauerman HL, Becker RC. Platelet functions beyond hemostasis. *Journal of thrombosis and haemostasis : JTH*. 2009; 7(11):1759–1766. [PubMed: 19691483]
105. Li C, Li J, Li Y, Lang S, Youghare I, Zhu G, Chen P, Ni H. Crosstalk between Platelets and the Immune System: Old Systems with New Discoveries. *Advances in hematology*. 2012; 2012:384685. [PubMed: 23008717]
106. Sabri A, Guo J, Elouardighi H, Darrow AL, Andrade-Gordon P, Steinberg SF. Mechanisms of protease-activated receptor-4 actions in cardiomyocytes. Role of Src tyrosine kinase. *The Journal of biological chemistry*. 2003; 278(13):11714–11720. [PubMed: 12522105]
107. Meune C, Spaulding C, Mahe I, Lebon P, Bergmann JF. Risks versus benefits of NSAIDs including aspirin in myocarditis: a review of the evidence from animal studies. *Drug safety : an international journal of medical toxicology and drug experience*. 2003; 26(13):975–981. [PubMed: 14583071]
108. Eysers S, Weatherall M, Shirtcliffe P, Perrin K, Beasley R. The effect on mortality of antipyretics in the treatment of influenza infection: systematic review and meta-analysis. *Journal of the Royal Society of Medicine*. 2010; 103(10):403–411. [PubMed: 20929891]
109. Shimazu T. A/H1N1 flu. Aspirin in the 1918 pandemic. *BMJ*. 2009; 338:b2398. [PubMed: 19528118]
110. Starko KM. Salicylates and pandemic influenza mortality, 1918–1919 pharmacology, pathology, and historic evidence. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. 2009; 49(9):1405–1410. [PubMed: 19788357]
111. Weitz JI, Gross PL. New oral anticoagulants: which one should my patient use? *Hematology / the Education Program of the American Society of Hematology American Society of Hematology Education Program*. 2012; 2012:536–540. [PubMed: 23233631]
112. Connolly S, Ezekowitz M, Brueckmann M, Clemens A, Yusuf S, Wallentin L. eLetter Re: Antoniak et al. PAR-1 contributes to the innate immune response during viral infection. *The Journal of clinical investigation*. 2013 <http://www.jci.org/eletters/view/66125#sec1>.
113. Malz R, Becher M, Westermann D, Schultheiss HP, Rauch U. Inhibition of coagulation factor Xa improves left ventricular function during CVB3-induced myocarditis [abstract]. Paper presented at the Clin Res Cardiol. 2012; 2012
114. Mason JW. Basic Research on Myocarditis: Superb but Unrequited. *Journal of the American College of Cardiology*. 2013
115. Lohman RJ, Cotterell AJ, Barry GD, Liu L, Suen JY, Vesey DA, Fairlie DP. An antagonist of human protease activated receptor-2 attenuates PAR2 signaling, macrophage activation, mast cell degranulation, and collagen-induced arthritis in rats. *The FASEB journal : official publication of the Federation of American Societies for Experimental Biology*. 2012; 26(7):2877–2887.
116. Sevigny LM, Zhang P, Bohm A, Lazarides K, Perides G, Covic L, Kuliopulos A. Interdicting protease-activated receptor-2-driven inflammation with cell-penetrating pepducins. *Proceedings of the National Academy of Sciences of the United States of America*. 2011; 108(20):8491–8496. [PubMed: 21536878]





**Figure. The thrombin PAR-1 pathway enhances TLR3-dependent innate immune responses in CVB3 myocarditis**

CVB3 infection leads to increased TF expression and activation of coagulation. Thrombin mediated PAR-1 cleavage contribute to TLR3 dependent IFN- $\beta$  due to increase in p38 activation on cardiac cells, such as cardiac fibroblasts. Additionally, PAR-1 can be activated by MMP13 and enhances TLR3 signaling during viral infection.

**Table 1**

Activating and inactivating proteases of PAR-1 and PAR-2

<b>Protease</b>	<b>Effect on PAR</b>	<b>Reference</b>
Thrombin	Primary activator of PAR-1	[59]
FXa	Activates PAR-1 and PAR-2	[60,61]
FVIIa	Activates PAR-2	[60,61]
MMP1 and MMP13	Activates PAR-1	[37,79,80]
Tryptase	Primary activator of PAR-2	[82]
Trypsin	Activates PAR-1 and PAR-2	[61]
Cathepsin G	Activates PAR-1	[72]
	PAR-2 inactivation on lung epithelial cells	[75]
Chymase	Reduces responses of fibroblasts to thrombin possible due to PAR-1 deactivation	[91]
	Activates PAR-2	[81]
Elastase	Activates PAR-1 on lung epithelial cells	[76]
	PAR-2 inactivation on lung epithelial cells	[75]
Papain	Induces PAR-2 dependent signaling in T-cells	[89]

**Table 2**

Roles of PAR-1, PAR-2 and different proteases in viral myocarditis

Manipulation	Observation	Reference
PAR-1 deficient mice	Increased CVB3 myocarditis due to reduced INF- $\beta$ responses	[37]
Overexpression of PAR-1 on cardiomyocytes <i>in vivo</i>	Reduced CVB3 myocarditis	[37]
PAR-2 deficient mice	Reduced CVB3 myocarditis	[95]
Thrombin inhibition with dabigatran etexilate in wild-type mice	Increased CVB3 myocarditis	[37]
MMP13 inhibition WAY170523) in wild-type mice	Increased CVB3 myocarditis	[37]
Fondaparinux (FXa inhibitor) in wild-type mice	Increased cardiac inflammation but improved heart function after CVB3 infection	[113]
Mast cell deficiency (loss of tryptase and chymase)	Reduced encephalomyocarditis virus myocarditis	[84]
Trypsin inhibition by aprotinin in wild-type mice	Reduced Influenza A virus myocarditis <i>in vivo</i> and <i>in vitro</i>	[67,68]
Neutrophil elastase inhibition by ZD0892 in wild-type mice	Reduced viral myocarditis by inhibiting immune cell infiltration	[77]
PAR-1 stimulation with TFLLR-NH(2)	Increased TLR3 dependent IFN- $\beta$ and CXCL10 expression in cardiac fibroblasts <i>in vitro</i>	[37]
PAR-1 inhibitor (SCH79797), MMP13 inhibitor WAY170523) and pan-MMP inhibitor (GM6001)	Reduced TLR3 dependent production of CXCL10 in cardiac fibroblasts <i>in vitro</i>	[37]
PAR-2 activation and PAR-2 overexpression	Reduced TLR3 dependent IFN- $\beta$ responses in cardiac fibroblasts and epithelial cells	[95,96]
PAR-2 inhibition by FSY-NH(2) or trypsin inhibition by aprotinin	Reduced influenza A virus replication in cardiomyocytes	[68]
Thrombin stimulation and PAR-1 expression	Increased susceptibility of endothelial cells to HSV	[92]
PAR-2 activation	Increased susceptibility of endothelial cells to HSV	[94]